

ASTRODYNAMICS SYMPOSIUM (C1)
Orbital Dynamics (2) (9)

Author: Dr. Josep-Maria Mondelo
Universitat Autònoma de Barcelona, Spain, jmm@mat.uab.cat

Dr. Àlex Haro
University of Barcelona, Spain, alex@maia.ub.es

COMPUTING QUASI-PERIODIC TRAJECTORIES OF THE RTBP IN $O(N \log N)$ OPERATIONS**Abstract**

Quasi-periodic (QP) trajectories play a fundamental role in many models in Astrodynamics, and provide many additional trajectory options in mission design beyond periodic motion [1,2]. They can be obtained locally by semi-analytical methods and globally by numerical ones. A numerical method that has progressively gained attention in the Astrodynamics community [1,2] is imposing invariance of a curve in the torus through the time- T flow (stroboscopic map), where T is one of the periods of the torus. Using a Fourier series approximation for the curve with N Fourier modes leads to a system of non-linear equations whose solution usually requires $O(N^3)$ operations. This is a computational bottleneck, especially for tori of more than two dimensions.

There is a new approach to the theoretical and computational study of invariant manifolds in dynamical systems known as the Parameterization Method [3,4,5]. In the computational side of this approach, differential correction is done in a functional framework, after the application of a transformation that makes use of geometrical properties that enable the computation of each Newton correction in a sequence of steps that are diagonal in either time or frequency domain. The computational benefit comes from the fact that the FFT allows to switch between these domains in $O(N \log N)$ operations.

In this paper, some theory and algorithms from [5] will be extended to enable the simultaneous computation of families of partially hyperbolic invariant tori together with the linear approximation of their stable and unstable manifolds. As a test, some of the families of tori of the RTBP (Lissajous, halo, quasi-halo) will be computed.

References

- [1] D. Guzzetti, N. Bosanac, A. Haapala, K. C. Howell, and D. C. Folta. Rapid trajectory design in the earth–moon ephemeris system via an interactive catalog of periodic and quasi-periodic orbits. *Acta Astronautica*, 126:439–455, 2016.
- [2] N. Baresi and D. J. Scheeres. Quasi-periodic invariant tori of time-periodic dynamical systems: applications to small body exploration. Paper IAC-16,C1,7,4,x32824, 67th International Astronautical Congress, 26-30 September 2016, Guadalajara, Mexico.
- [3] R. de la Llave, A. González, À. Jorba, and J. Villanueva. KAM theory without action-angle variables. *Nonlinearity*, 18(2):855–895, 2005.
- [4] À. Haro and R. de la Llave. A parameterization method for the computation of invariant tori and their whiskers in quasi-periodic maps: numerical algorithms. *Discrete Contin. Dyn. Syst. Ser. B*, 6(6):1261–1300, 2006.
- [5] A. Haro, M. Canadell, J.-L. Figueras, A. Luque, and J. M. Mondelo. *The parameterization method for invariant manifolds: from rigorous results to effective computations*. Applied Mathematical Sciences. Springer, 2016.