

45th IAA SYMPOSIUM ON THE SEARCH FOR EXTRATERRESTRIAL INTELLIGENCE (SETI) –
The Next Steps (A4)
SETI 1: SETI Science and Technology (1)

Author: Dr. Alexander Panov
Skobeltsyn Institute of Nuclear Physics, Russian Federation, panovenator@gmail.com

Dr. Nikolaj Bursov
Russian Academy of Sciences, Russian Federation, nmb@sao.ru

Mrs. Liliya Filippova
Russian Federation, lfilippova@yandex.ru

Dr. Lev Gindilis
Sternberg Astronomical Institute of Moscow Lomonosov State University, Russian Federation,
lgindilis@mail.ru

Dr. Claudio Maccone
International Academy of Astronautics (IAA) and Istituto Nazionale di Astrofisica (INAF), Italy,
clmaccon@libero.it

Mr. Vladimir Philippov
Russian Federation, vvfopal@gmail.com

Mr. Evgeny Starikov
Russian Federation, e.starikoff@gmail.com

Mr. James Wilson
Australia, eagleny@gmail.com

SETI OBSERVATION WITH THE RATAN-600 TELESCOPE IN 2015 YEAR AND DETECTION OF A
STRONG SIGNAL IN THE DIRECTION OF HD164595

Abstract

39 targets (37 stars and 2 globular clusters) were observed within SETI program with RATAN-600 telescope (Russia) in 2015. The observations were carried out at 6.2 cm, 2.7 cm, 1.4 cm and 1.0 cm wavelengths in broad bandwidths. The sensitivity of the telescope is determined by the noise temperature of the receiver being about 5 mK. The mode of observation was transition of object through a fixed telescope aperture. The duration of one transition was about 2 sec and 6017 separate patterns of transitions of targets were recorded during 2015. A strong signal had been received on 2.7 cm wavelength exactly during transition of the star HD164595 May 15, 2015. The probability of a casual noise simulation of a signal was estimated as one per more than 6 years of observations with 6000 separate observations a year. HD164595 should be considered as a good candidate for SETI observations.