IAF/IAA SPACE LIFE SCIENCES SYMPOSIUM (A1) Astrobiology and Exploration (6)

Author: Dr. Laurence Lemelle CNRS, France

Dr. Christophe Place **CNRS**, France Mrs. Eleonore Mottin **CNRS**, France Mr. Denis LeTourneau **CNRS**, France Dr. Guillaume Nonglaton Commissariat à l'énergie atomique et aux énergies alternatives (CEA), France Mr. Pierre Marcoux Commissariat à l'énergie atomique et aux énergies alternatives (CEA), France Dr. Jeremie Teisseire **CNRS**. France Mr. Emmanuel Garre **CNRS**. France Dr. Lucie Campagnolo MEDES, France Mrs. Cecile Thevenot MEDES, France Dr. Alain Maillet MEDES - IMPS, France Mr. Sebastien Rouquette CADMOS-CNES, France Mr. Patrice Benarroche CADMOS-CNES, France Mr. Sebastien Barde

A MATISS-1 OPTICAL SURVEY OF THE PARTICLES CLASSES CONTAMINATING VARIOUS SURFACES IN THE ISS

CADMOS-CNES, France

Abstract

Biocontamination in manned spacecrafts and in future habitats could have significant impacts on crew health and biodegradation of equipment. In particular, there is a need for decreasing the dependency of crew action for cleaning actions. Anti-microbial surfaces that inhibit or reduce the ability of microorganisms to grow on the surfaces are of high interest. In the context of future manned mission scenarios of longer duration, higher isolation and the utilization of an increasing number of closed-loop life support systems, the coating or surface modification must be effective 'in-use'.

The MATISS experiment (Microbial Aerosol Tethering on Innovative Surfaces in the International Space Station) aims to demonstrate that surfaces with hydrophobic properties already implemented in

numerous industrial fields could be a possible answer applicable on the scale of spacecraft. By reducing the contact area of the droplets of water with surfaces, the hydrophobicity allows at the first order to limit the fraction of the surface which is contaminated. It also limits the adhesion of microorganisms to surfaces. In this respect, a spatialized set-up was developed.

A MATISS set-up that protects the glass surfaces while sampling ISS atmosphere was conceived and mounted glass lamella which surfaces were modified to become hydrophobic applying industrializable processes controlled by two RD French groups: the CEA Léti and Saint Gobain. 4 sample holders were exposed 6 months in the Columbus module and returned in their ziplock bags for analyses under confinement in laboratories. PCR was carried out on the biocontaminations sampled on the external surface of the holders. Systematic imaging of the surfaces under confinement was carried out at low and high magnification.

An outcome of the optical survey of the exposed surface is the indexation of 5500 particles (average density <2 particles / mm2). Based on their sizes and shapes, we can distinguish 5 classes of particles: textile fibers (Area>1500 μ m2), irregular and angular patches (50 μ m2<Area<1500 μ m2), and very much smaller particles consistent with colonies (4.5 μ m2 < Area < 15 μ m2) and cocci and rod-shaped particles (0.5 μ m2 < Area < 4.5 μ m2). The influence of the surface coating on the distribution of each class of particle displays a clear impact of the FDTS coating on the fibers and the smallest particles distributions that will be more systematically investigated from the on-going experiments MATISS-2 and 2.5.