

IAF/IAA SPACE LIFE SCIENCES SYMPOSIUM (A1)
Medical Care for Humans in Space (3)

Author: Ms. Xinyu Wang

Shanghai Engineering Center for Microsatellites, Chinese Academy of Sciences (CAS), China,
luckygirlwxy@163.com

Dr. Yonghe Zhang

Shanghai Engineering Center for Microsatellites, Chinese Academy of Sciences (CAS), China,
zhangyh@microsate.com

Dr. Guopeng Ding

Shanghai Engineering Center for Microsatellites, Chinese Academy of Sciences (CAS), China,
dianxin0307@163.com

Prof. Ping Hu

Shanghai institute for Biological Sciences, Chinese Academy of Sciences (CAS), China, hup@sibcb.ac.cn

Ms. Wenxiu Zhang

Shanghai Engineering Center for Microsatellites, Chinese Academy of Sciences (CAS), China,
zwxy@mail.ustc.edu.cn

Mr. Ming Guo

Shanghai Engineering Center for Microsatellites, Chinese Academy of Sciences (CAS), China,
googlm@163.com

Dr. Zhongguang Yang

Shanghai Engineering Center for Microsatellites, Chinese Academy of Sciences (CAS), China,
yangzg@microsate.comMEASUREMENT SYSTEM OF MYOTUBE MORPHOLOGICAL PARAMETERS FOR SPACE LIFE
EXPERIMENT MISSION ABOUT MUSCLE ATROPHY IN ASTRONAUTS**Abstract**

In long-term space life, muscle atrophy in astronauts has still confused us, which may be caused by microgravity, space radiation or other unknown factors. Therefore, a space life experiment will be carried out by Shanghai Engineering Center for Microsatellites (SECM) to explore the underlying molecular mechanisms of space muscle atrophy via cultured muscle stem cells and the differentiated myotubes. In this model, detecting morphological parameters of myotubes is widely used for evaluating the severity of muscle atrophy. However, it is hard to perform the measurement precisely with bright-field microscopy images, which is the most common way to monitor the muscle differentiation process and the extend of muscle atrophy. In these images, many myotubes are minute and similar to the backgrounds. Moreover, many myotubes are overlapping with each other and it is difficult to distinguish one myotube from others. To our knowledge, existing myotube morphological measurement is carried out manually by scientists via software, such as Image-Pro Plus. It highly depends on the experience of operators, which can not be implemented in the space environment. In this paper, we present an automatic myotube morphological parameter measurement system, which can measure the area, the width and the length of myotubes. Due to the complexity of myotube images, the deep learning method is adapted to achieve the high accuracy instance-aware segmentation. We adopt U-net deep learning framework to segment myotube images and then extract contours of myotubes to measure the morphological parameters based on curvature analysis. Using a few of manually-annotated fluorescent images for pre-training and 400 manually-annotated

microscopy images for training, the neural network can generate high accuracy myotube segmentation masks that allow us to analyze the morphology of myotubes. The number of pixels in each segmentation mask is regarded as the myotube area. And then, we detect the cytoskeleton based on the curvatures of each segmentation mask contour to compute the myotube width and length. Comparing with the manual measurement, our approach is more objective and saves a lot of time.